



A Cross-sectional Study of Psychosocial Problems Following Therapeutic Abortion With the Mother's Spiritual Experiences

Sedighe Alipanahpour¹, Mahnaz Zarshenas², Mina Taheri², Marzieh Akbarzadeh^{3*}

Abstract

Objectives: Abortion can be stressful for the family and may lead to psychological problems. The question arises whether religious attitudes can be restructured into women who experience induced and spontaneous abortions. In this regard, the present study aimed to investigate the relationship of post-traumatic stress disorder (PTSD) after therapeutic abortion (Induced and spontaneous) with the mother's spiritual experiences.

Materials and Methods: This cross-sectional study with a sample size of 104 people was conducted in 2018-2019 in the selected hospitals of Shiraz University of Medical Sciences. The convenience sampling method was used in 2018. Research tools including the Mississippi PTSD Scale questionnaire were implemented for measuring stress and religious attitudes and completed immediately and a month after abortion. Data were analyzed by SPSS software using one-way ANOVA, least significant difference post hoc test, and paired t-test or Wilcoxon test.

Results: Based on the results, 68.7%, 71.8%, and 72.7% of the abortion group with forensic medical letter, other causes, and spontaneous abortion had a high level of religious attitudes, respectively. In addition, 78.1%, 69.2%, and 72.7% of those who had an abortion with a forensic medical letter, underwent abortion for other causes, and experienced a spontaneous abortion had moderate PTSD immediately after abortion, respectively. Further, 62.5%, 64.1%, and 66.7% of women having an abortion with forensic medical letter, undergoing abortion for other causes, and experiencing spontaneous abortion had moderate PTSD one month after abortion, respectively. The findings revealed no significant relationship between PTSD differences immediately and a month after the abortion in the subjects ($P=0.175$).

Conclusions: The research community had a high religious attitude while having no association with PTSD reductions. More than half of the pregnancies were unwanted, which may be due to stress levels.

Keywords: Abortion, Post-traumatic stress, Religious attitude, Mothers, Spontaneous, Induction, Forensics

Introduction

Terminology and Prevalence

Pregnancy loss can take many forms including abortion, stillbirth, and miscarriage and is accompanied by maternal and familial stress in all these cases. Spontaneous abortion is the most common and serious complication in early pregnancy, occurring in 17%-22% of all known pregnancies (1). According to the Centers for Disease Control and the World Health Organization, abortion is defined as the termination of a pregnancy before the 20th week of pregnancy or the birth of a fetus weighing less than 500 grams (2). Its prevalence in the United States is 15% of the known pregnancies with spontaneous abortion and about one-third of pregnancies of women with selective abortion (3). As mentioned by Sedgh et al (4), Dakar, the capital of Senegal, had 21 abortions per 1000 to 44-400 women, which was higher than those in other parts of the country (16 per 1000). In one study in Iran, the prevalence

of abortion was 8.3% (5) and in another study in Tehran, this value was estimated as 8.7% of those among married women ending in abortion, namely, in every 100 known pregnancies (6).

The Type of Traumatic Reactions Following Abortion Experiences

Hope for the future, feeling of satisfaction, and early bonding with the unborn child are among the complex emotional responses to pregnancy due to its effective nature. On the other hand, abortion can be stressful for family members, doctors, and others in the social support system (7). Some women experience anxiety, anger, post-traumatic stress, and feeling of guilt about childbearing in the future (8) and its detrimental effects on the couple's relationship (9). According to Langkaas et al (10), such feelings can be part of a persistent, overactive, or functional disorder that is associated with post-traumatic

Received 5 August 2020, Accepted 23 January 2021, Available online 4 June 2021

¹Department of Midwifery, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran. ²Community-Based Psychiatric Care Research Center, Department of Midwifery, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran. ³Department of Midwifery, Maternal-fetal medicine Research Center, School of Nursing and Midwifery, Shiraz University of Medical Sciences, Shiraz, Iran.

*Corresponding Author: Marzieh Akbarzadeh, Tel: +98 0711-6474250, Fax: +98 0711-647425, Email: akbarzadm@sums.ac.ir



Key Messages

- ▶ Women's psychological responses to abortion and PTSD are influenced by complex socio-cultural and ethnic factors, and psychological problems may persist for a long time after physical recover.
- ▶ The existence of religious beliefs as an effective factor in the prevention and treatment of mental disorders, if used effectively, may affect the ability of individuals to adapt and cope with the crisis of abortion. Positive hope and attitude will be effective in reducing PTSD.

stress disorder (PTSD). Abortion is associated with moderate to severe mental, social, and occupational risk including psychological problems such as depression and anxiety, suicidal behaviors, PTSD, alcohol or marijuana use, and smoking (7,8,11). A cohort study in Sweden was performed on 1457 women with induced abortion. The prevalence of PTSD before and after abortion was 4.3% and 23.5%, respectively, and there was a high level of anxiety, depression, and dropout. Evidence showed that within one month after abortion, 10% of women had acute stress disorder and 1% had PTSD (7). Other studies reported that people respond to the bitter experience of abortion in the form of fear and helplessness, constantly imagining the event in their minds. The possibility of post-traumatic stress increases as well. However, no study has focused on religious attitudes and post-traumatic stress in recent years (12-15).

Religious Beliefs and Practices Defined as a Method of Coping

Many scholars consider having religious beliefs as an influential factor in people's health. This factor can also be effectively used in the treatment and prevention of mental disorders and can increase the ability of individuals to adapt to and deal with illnesses (16). From the mental health point of view, religion offers many guidelines that can help people in creating principles for their lives. Religious beliefs can easily make the pressures, stresses, and uncertainties of life tolerable (17). In addition, religion is a source of support for individuals facing problems, and religious beliefs and practices are defined as a method of coping that uses religious resources such as prayer, trust, and the appeal to God. Religion further leads to the mental health of the individual and society through various mechanisms. Moreover, performing religious behaviors such as prayer, honesty, and faith in God and reading religious books also create inner peace by providing hope and encouragement regarding finding a positive view of the situation. Other mechanisms include having religious mental health, hope, and motivation and positivity in religion, creating an emotional and social support network, giving a clear and decisive answer to the concept of creation, world, and life, along with defining

suffering, pain, and deprivation (17,18).

Numerous studies have shown not only a positive relationship between religion and mental health but also the positive effects of religious beliefs and practice on one's mental and physical health (19). Abortion is a controversial issue in all religions and medical societies and schools. From the Islamic point of view, criminal abortion is a sin and unacceptable except for medical cases, especially in certain circumstances. It should be noted that the majority of people are Muslim in Iran. From the perspective of Shiite jurisprudence in Islam with reference to the Holy Quran, all human beings, including the fetus, are equal in the principles of "human dignity" and "right to life". In general, from the perspective of Shiite jurists, abortion is forbidden and punished under the Islamic penal code. Most Shiite jurists allow abortion only in the case of unhealthy and incomplete fetuses and highly urgent medical cases (i.e., heart disease, hypertension, and the like). Additionally, abortion can be performed with the permission of judicial authorities and a forensic medical certificate in cases that the conditions exacerbate the effects of the disease and threaten the mother's life (20-23).

Necessity and Reasons for Conducting Research

Given the high incidence of abortion and its subsequent physical and psychological complications, some studies demonstrated that God-given healing and faith-based therapy are well-organized, psychological-based methods of high importance in treating diseases and alleviating induced pain, anxiety, depression, and tensions. In this regard, the current study mainly sought to determine whether the level of women's religious beliefs in society can be considered as a factor in promoting the attitude toward abortion. More precisely, whether it is possible to expect a more favorable attitude toward abortion by increasing the level of religious belief or whether there is a correlation between the level of women's religious attitudes in PTSD and the types of abortions, and how it is effective in PTSD. Therefore, the study focused on evaluating the relationship between religious attitudes and PTSD in all types of spontaneous and induced abortion permitted by forensic medicine in the hospitals affiliated to Shiraz University of Medical Sciences in 2018 in order to use the results to help a large group of pregnant women.

Materials and Methods

This cross-sectional study was performed in the selected hospitals of Shiraz University of Medical Sciences (i.e., Hazrat Zeinab, Shahid Faghihi, Hafez, and Shoostari hospitals) due to their availability and a high number of referrals in 2018-2019. The sample size was calculated as 82 patients based on the correlation coefficient estimated from the study by Cowchock et al (24) with a loss probability of 10%. In general, 104 subjects were enrolled given the mothers' willingness to participate in the study.

The inclusion criteria were being 10-49 years old, Iranian, and literate and having experience of one type of abortion while not suffering from chronic illness (i.e., heart disease, hypertension, and diabetes) or using any psychological treatment including medication and psychotherapy. The types of abortion were induced abortion with a forensic medical letter due to fetal abnormalities or maternal complications, induced abortions with other medical causes (e.g., blighted ovum), and spontaneous abortion occurring without any intervention. On the other hand, the exclusion criteria were women's willingness to withdraw from the study at any time and any other adverse events in life that can affect the mother's grief (e.g., financial crisis or bereavement).

A simple purposive sampling method was used in this study, and the sampling lasted for 4 months from September 2018 to January 2019. General information questionnaire and medical and midwifery history and type of abortion until the moment of abortion were recorded after obtaining written consent from eligible individuals. Then, Mississippi PTSD Scale was applied to measure stress levels, and a religious attitude questionnaire was completed for all the samples. The questionnaires were completed immediately after the abortion and one month later via phone calls.

Study Design

After obtaining the code of ethics from the Ethics Committee of Shiraz University of Medical Sciences, the researcher was introduced to the hospitals. Then, the researcher referred to three hospitals. The target population consisted of all women candidates for abortion for any reason. They were given a verbal lecture and written information about the study objectives, approach, and necessary training on how to fill out the questionnaires, and were assured of data confidentiality. Willing women signed a written consent form and participated in the study. They filled out the general information questionnaire, and their medical history and midwifery records were recorded until the time of the abortion including the type of abortion, causes and complications, and type of treatment. Next, a standardized questionnaire and the recovery assessment scale-revised were applied to measure stress levels and religious attitudes, respectively. Four weeks after the abortion, subjects were asked to re-complete the stress and religious questionnaires. It should be noted that the information about the causes, complications, and type of treatment is not included in this article.

Data Collection Tools

a. The demographic questionnaire consists of 60 researcher-made items in two sections of demographic (14 items) and midwifery (46 items) information. To determine validity, the final version of the questionnaire approved by the research team

was given to 10 expert professors of the schools of nursing and midwifery of Shiraz Universities of Medical Sciences. The validity of the questionnaire was confirmed after the application of the professors' recommendations.

- b. Mississippi PTSD Scale includes 35 items scored from 1 to 5. The total score ranges from 35 to 175 with scores of 107 and above indicating a person with PTSD (25-29). This scale has been validated in Iran by Goodarzi and with a Cronbach's alpha coefficient of 0.92 (28). Three tools of life events inventory, the PTSD index, and the Padua inventory were used to determine the concurrent validity of this scale. The correlation coefficients of the Mississippi scale with each tool were reported as 0.23, 0.82, and 0.75, respectively (29). This validation process was utilized as a basis for the present study.
- c. The Religious Attitude Questionnaire consists of 25 items scored based on a Likert-type scale ranging from 1 to 5, classifying the participants into three levels of high (scored 100 and above), low (scored 50 and lower), and moderate (scored 51-99) religious attitudes (19,30,31). Its Cronbach's alpha coefficient was estimated as 0.954. The reliability and validity indexes reported by Ebrahimi et al were the basis of the present study (30).

Statistical Analysis Methods

One-way ANOVA, least significant difference, and post hoc tests were used to compare stress levels in the three groups. Paired *t* test or Wilcoxon test was also used to compare stress in each group. In addition, one-way ANOVA was applied to express differences in scores immediately and one month after abortion in addition to comparing the three groups according to their religious attitudes.

Results

The mean age of the subjects was 30.62 years, of whom 89.4% were housewives and 41.3% had a high school diploma. The mean age of their spouses was 34.94%, of whom 77.8% were self-employed and 35.6% had a high school diploma (Table 1).

In the spontaneous abortion group, 72.7% and 28.8% had high and medium levels of religious attitudes, respectively (Table 2). The induced abortion group with a forensic medical letter, the induced abortion group with other causes, and the spontaneous abortion group had a PTSD score of 30.8%, 37.5%, and 31.7%, respectively, and the highest stress scores in all three groups were moderate (Table 3).

One month after abortion, 62.5%, 64.1%, and 66.7% of subjects in the induced abortion group with a forensic medical letter, the induced abortion group with other causes, and the spontaneous abortion group had moderate PTSD, respectively (Table 4).

Table 1. Demographic Characteristics of the Study Subjects

Variable	Component	Induced Abortions (Forensic Medicine)	Induced Abortions (Other Etiology)	Spontaneous Abortions Spontaneous Abortions	Total	
		n (%)	n (%)	n (%)	n (%)	
Job	Mother	Housewife	29 (90.6)	35 (89.7)	29 (87.9)	93 (89.4)
		Self-employment	1 (3.1)	1 (2.6)	1 (3)	3 (2.9)
		Employment	2 (6.3)	3 (7.7)	3 (9.1)	8 (7.7)
	Father	Unemployment	1 (3.1)	0 (0)	0	1 (1)
		Self-employment	23 (71.9)	32 (82.1)	26 (78.8)	81 (77.8)
Education	Mother	Employment	8 (25)	7 (17.9)	7 (21.2)	22 (21.2)
		Elementary	9 (28.1)	5 (12.8)	3 (9.1)	17 (16.4)
		Middle school	7 (21.9)	6 (15.4)	6 (18.2)	19 (18.3)
	Father	High School (Diploma)	9 (28.1)	18 (46.2)	16 (48.5)	43 (41.3)
		Licensee and postgraduate	7 (21.9)	10 (25.6)	8 (24.2)	25 (24.0)
		Elementary	4 (12.5)	4 (10.2)	3 (9.1)	11 (10.6)
	Father	Middle school	11 (34.4)	12 (30.8)	12 (36.4)	35 (33.7)
		High school (Diploma)	11 (34.4)	15 (38.5)	11 (33.3)	37 (35.6)
		Licensee and Postgraduate	6 (18.7)	8 (20.5)	7 (21.2)	21 (20.1)
Age	Mother	20 and lower	1 (3.1)	0 (0)	3 (9.1)	4 (3.8)
		21-30	10 (31.3)	20 (51.3)	17 (51.5)	47 (45.2)
		31-40	16 (50)	18 (46.1)	13 (39.4)	47 (45.2)
		41 and higher	5 (15.4)	1 (2.6)	0 (0)	6 (5.8)
	Father	Mean	32.53	30.51	28.88	30.62
		20-30	5 (15.6)	5 (12.8)	12 (36.4)	22 (21.15)
		31-40	19 (59.4)	30 (76.9)	17 (51.5)	66 (63.46)
		41 and higher	8 (25)	4 (10.5)	4 (12.1)	16 (15.4)
		Mean	36.6	35.12	33.12	34.94
Total		32 (100)	39 (100)	33	104 (100)	

Table 2. The Level of Religious Attitudes in the Induced and Spontaneous Abortion Groups

RA Score	Induced Abortions (Forensic Medicine)	Induced Abortions (Other Etiology)	Spontaneous Abortions	Total
	n (%)	n (%)	n (%)	n (%)
Medium (50-100)	10 (31.3)	11 (28.2)	9 (27.3)	30 (28.8)
High (100-125)	22 (68.7)	28 (71.8)	24 (72.7)	74 (71.2)
Total	32 (100)	39 (100)	33 (100)	104 (100)

RA: Religious attitude score.

Table 3. Comparison of PTSD Intensity in the Induced and Spontaneous Abortion Groups Immediately After Abortion

PTS Score	Induced Abortions (Forensic Medicine)	Induced Abortions (Other Etiology)	Spontaneous Abortions	Total
	n (%)	n (%)	n (%)	n (%)
Low (35-70)	6 (18.8)	7 (17.9)	7 (21.2)	20 (19.2)
Medium (70-107)	25 (78.1)	27 (69.2)	24 (72.7)	76 (73.1)
High (107-175)	1 (3.1)	5 (12.8)	2 (6.1)	8 (7.7)
Total	32 (30.8)	39 (37.5)	33 (131.73)	104 (100)

PTS: Post-traumatic stress score; PTSD: Post-traumatic stress disorder.

One-way ANOVA was used to compare the three groups in terms of religious attitudes. The results showed that there was no statistically significant difference between the three groups regarding religious attitudes (Table 5). Spearman correlation was applied to determine the correlation between PTSD score changes immediately and one month after abortion with the religious attitude in general and in each of the three groups. Based on the results, the changes in the scores were not significant. In other words, the difference in PTSD was not significant immediately and one month after the abortion ($P = 0.175$).

Data analysis on the relationship between the PTSD score and religious attitudes revealed that the difference in PTSD scores immediately and one month after abortion was significant and inverse only in the case of induced abortion with other causes ($P = 0.027$) while not being statistically significant in other cases. In other words, higher scores of the religious attitudes led to lower PTSD scores before and after abortion (Tables 5 and 6).

Discussion

Our results demonstrated that patients with abortion

Table 4. Comparison of PTSD Intensity in the Induced and Spontaneous Abortion Groups a Month After Abortion

PTS Score	Induced Abortions (Forensic Medicine)	Induced Abortions (Other Etiology)	Spontaneous Abortions	Total
	n (%)	n (%)	n (%)	n (%)
Low (35-70)	11 (34.4)	12 (30.8)	11 (33.3)	34 (32.7)
Medium (70-107)	20 (62.5)	25 (64.1)	22 (66.7)	67 (64.3)
High (107-175)	1 (3.1)	2 (5.1)	0 (0)	3 (2.9)
Total	32 (100)	39 (100)	33 (100)	104 (100)

PTS: Post-traumatic stress score; PTSD: Post-traumatic stress disorder.

Table 5. Comparison of Religious Attitudes in the Induced and Spontaneous Abortion Groups

Induced Abortions (Forensic Medicine)	Induced Abortions (Other Etiology)	Spontaneous Abortions	Index	P Value
8.2±103.1	9.7±102.2	9.6±103.4	0.168	0.848

Table 6. The Relationship Between the PTSD Score and Religious Attitudes

PTS Score	Induced Abortions (Forensic Medicine)		Induced Abortions (Other Etiology)		Spontaneous Abortions		Total	
	P Value	r	P Value	r	P Value	r	P Value	r
Immediately after abortion	0.951	-0.02	0.415	0.136	0.802	-0.04	0.847	0.01
A month after abortion	0.768	-0.05	0.325	-0.164	0.272	-0.11	0.212	-0.12
Variation	0.747	0.06	0.027	-0.259	0.740	-0.05	0.175	-0.134

PTSD: Post-traumatic stress disorder.

had a high religious attitude. The results of the group with induced abortions with other causes showed that the difference in stress immediately after an abortion and a month later was significant and had an inverse relationship ($P = 0.027$), meaning that the higher was the religious attitude score, the lower was the stress score. A review study on the role of mothers' religious beliefs in dealing with pregnancy loss represented that religion has a significant impact on the parents' acceptance of fetal loss and their recovery from such tragic events (32). Based on the results of the study by Layer et al on the effect of spiritual group intervention on post-abortion sadness and mourning, PTSD symptoms were less responsive to intensive interventions (33). Likewise, Cowchock et al evaluated the effect of religious beliefs on post-pregnancy sadness and found that belief in God reduced sadness and grief scores 4-6 weeks after fetal loss (24,33). All the above-mentioned studies showed that higher religious attitudes and spiritual beliefs reduced PTSD in mothers with induced abortion. These results are consistent with those of our study.

The impact of one's spiritual beliefs on the interpretation of events can facilitate the process of accepting events. In other words, spirituality is described as an umbrella that covers various concepts such as spiritual health, beliefs, and spiritual adjustment (34). Religion satisfies one's basic needs and fills moral, emotional, and spiritual gaps (35). Furthermore, religious beliefs and behaviors help the individuals cope with stress, create hope and positive attitudes, strengthen their inner peace, and successfully cope with stress (32,36-38). They also have better mental health and less depression and anxiety (19,31,39-42).

Our results demonstrated that the majority of abortion

patients with a forensic medical letter and spontaneous abortion had religious attitudes. There was no significant relationship between PTSD and religious attitudes in the forensic medicine group.

Some studies reported that traumatic events cause PTSD. However, the severity of the initial response and the number of people showing these reactions markedly decreased over time (43-45). A study in the US evaluated the mental health of women who had unwanted pregnancies and abortions one hour later, one month later, and two years later and concluded that most of these women did not experience psychological problems (46). The results of these studies are in line with the findings of our study.

Some studies showed that spontaneous abortion does not lead to psychological complications although factors such as pre-abortion mental health, domestic violence, pregnancy tendency, and economic status affect depression and PTSD (47,48). However, induced abortion had some negative psychological effects on women because they have been deprived of a healthy child. Based on the results of Allahdadian and Irajpou, religion and religious beliefs reduce the sadness and stress caused by fetal loss and prepare the parents to accept it (32).

On the other hand, other studies indicated that no specific psychological changes occur after an abortion, and as a result, there is no relationship between abortion and post-abortion mental health. The problems that occur after abortion are due to violence and social deprivation in these women before the abortion. When these factors were taken into account during the analysis of the PTSD data, abortion was not associated with subsequent psychological problems (49-51).

In this study, 53.8% of the pregnancies were unwanted (50% of these women did not use a reliable contraceptive method), and they did not take any action for abortion and considered spontaneous abortion or the need for induced abortion as a kind of a divine blessing because of their religious beliefs and attitudes about their feeling of guilt about abortion. They were wholeheartedly happy about the incident. Therefore, the unwillingness of these women to become pregnant was one of the reasons for this issue and the lack of a decrease in the stress level of these people despite their high religious attitudes. Otherwise, the authors can have no specific justification in this regard. In addition, permitted abortions by forensic medicine usually have fetal problems or abnormalities, or the mother has a serious illness that can lead to a life-threatening pregnancy.

The role of religion in dealing with stressors is highly complex and not fully understood yet (52). In a review study, Becker et al found that religious beliefs had positive effects on the relief of sadness (53). A constant heart-to-heart connection with God and friendship with Him and His remembrance brings reassurance to the heart and relieves all anxiety and worries of human existence. Great people, with a big and calm heart, spend their lives with security and pride. Verse 62 of *Surah Yunus* in the Holy Quran says:

“Remember, there is neither fear nor regret for the friends of God.”

The same belief in God is also mentioned in many other verses (*al-Ahaqaf* 13, and *al-Fater* 24).

Some evidence suggests that there may be a beneficial relationship between religion and mental peace. Some people may turn to God by losing something while others may turn away from God (54). Anxiety and depression have long been known to be associated with abortion. Several studies have confirmed a relationship between abortion experience and the onset of PTSD (55-57). The psychological responses of women to abortion and PTSD symptoms are influenced by complex sociocultural and ethnic factors (58,59). Accordingly, the following recommendations are presented to reduce the incidence of abortion and PTSD.

1. Given the religious context in Iran, attention should be focused on primary prevention. In this respect, it is recommended that authorities set up services for all members of the community, both single and married. Access to necessary services should be available 24/7.
2. It is recommended that the existing abortion policies should be revised with an emphasis on maternal health. Of course, the new fatwas of religious scholars and laws that make abortion eligible in the areas of hardship in the period before the insufflation of spirit could pave the way for the revision of laws by courts and lawmakers to reduce or even avoid unsafe abortions.

Conclusions

The results of our study showed that induced abortions (with other etiologies) had an inverse relationship with religious attitude and there was no significant statistical relationship with religious attitudes in the case of spontaneous and induced (Forensic Medicine) abortions.

In this study, 53.8% of the pregnancies were reported as unwanted pregnancy cases, and 36.8% of women took no action for abortion due to religious beliefs and attitudes about the guilt of abortion despite the parents' unwillingness to keep the unwanted fetus. Therefore, they considered spontaneous abortion or the need for induced abortion as a divine gift. It can be argued that their unwanted pregnancy was one of the reasons for the lack of a decrease in stress levels in these women. Some studies demonstrated that although no specific psychological changes occur after an abortion, some factors such as pre-abortion mental health, domestic violence, pregnancy tendency, and economic status contribute to depression and PTSD. However, induced abortion has had some negative psychological effects on women because women have been deprived of a healthy child. It seems that further studies are needed to confirm or disprove the existence of this relationship. If there is a relationship, the religious trainings by midwives and other health care providers should be used to help reduce PTSD in women with abortion.

Authors' Contribution

MA and SA prepared the first draft of the manuscript and MA and MZ made critical revisions to the paper and responded to the reviewers. MT and MA helped the Surge Articles.

Conflict of Interests

The authors declare no conflict of interest, financial or otherwise.

Ethical Issues

This study was conducted with the code number of IR.SUMS.REC.1396.S673 and financially supported by Shiraz University of Medical Sciences. Subjects were free in providing their details (questionnaires were marked with codes and at each stage) and could leave the study at any stage. The researchers sought to ensure that all participants' rights were respected following the Helsinki Convention on Ethics.

Financial Support

Vice-chancellor of Shiraz University of Medical Sciences, Shiraz, Iran.

References

1. Grippo A, Zhang J, Chu L, et al. Air pollution exposure during pregnancy and spontaneous abortion and stillbirth. *Rev Environ Health*. 2018;33(3):247-264. doi:10.1515/reveh-2017-0033
2. Cameron MJ, Penney GC. Terminology in early pregnancy loss: what women hear and what clinicians write. *J Fam Plann Reprod Health Care*. 2005;31(4):313-314. doi:10.1783/147118905774480761
3. Hamama L, Rauch SA, Sperlich M, Defever E, Seng JS.

- Previous experience of spontaneous or elective abortion and risk for posttraumatic stress and depression during subsequent pregnancy. *Depress Anxiety*. 2010;27(8):699-707. doi:10.1002/da.20714
4. Sedgh G, Sylla AH, Philbin J, Keogh S, Ndiaye S. Estimates of the incidence of induced abortion and consequences of unsafe abortion in Senegal. *Int Perspect Sex Reprod Health*. 2015;41(1):11-19. doi:10.1363/4101115
 5. Motavalli R, Alizadeh L, Namadi Vosoughi M, Shahbazzadegan S. Evaluation of the prevalence, reasons and consequences of induced abortion in women of Ardabil in 2011. *J Ardabil Univ Med Sci*. 2012;12(4):384-391. [Persian].
 6. Erfani A. Induced abortion in Tehran, Iran: estimated rates and correlates. *Int Perspect Sex Reprod Health*. 2011;37(3):134-142. doi:10.1363/3713411
 7. Bowles SV, James LC, Solursh DS, Yancey MK, Epperly TD, Folen RA, et al. Acute and post-traumatic stress disorder after spontaneous abortion. *Am Fam Physician*. 2000;61(6):1689-1696.
 8. Haghparast E, Faramarzi M, Hassanzadeh R. Psychiatric symptoms and pregnancy distress in subsequent pregnancy after spontaneous abortion history. *Pak J Med Sci*. 2016;32(5):1097-1101. doi:10.12669/pjms.325.10909
 9. Hutti MH, Armstrong DS, Myers JA, Hall LA. Grief intensity, psychological well-being, and the intimate partner relationship in the subsequent pregnancy after a perinatal loss. *J Obstet Gynecol Neonatal Nurs*. 2015;44(1):42-50. doi:10.1111/1552-6909.12539
 10. Langkaas TF, Hoffart A, Øktedalen T, Ulvenes PG, Hembree EA, Smucker M. Exposure and non-fear emotions: a randomized controlled study of exposure-based and rescripting-based imagery in PTSD treatment. *Behav Res Ther*. 2017;97:33-42. doi:10.1016/j.brat.2017.06.007
 11. Kulathilaka S, Hanwell R, de Silva VA. Depressive disorder and grief following spontaneous abortion. *BMC Psychiatry*. 2016;16:100. doi:10.1186/s12888-016-0812-y
 12. Schwandt HM, Creanga AA, Danso KA, Adanu RM, Agbenyega T, Hindin MJ. A comparison of women with induced abortion, spontaneous abortion and ectopic pregnancy in Ghana. *Contraception*. 2011;84(1):87-93. doi:10.1016/j.contraception.2010.10.011
 13. Wallin Lundell I, Sundström Poromaa I, Frans O, et al. The prevalence of posttraumatic stress among women requesting induced abortion. *Eur J Contracept Reprod Health Care*. 2013;18(6):480-488. doi:10.3109/13625187.2013.828030
 14. Alekseeva N, Horton R, Geller F, McGee J, Minagar A. Psychiatric disorders and pregnancy. In: Minagar A, ed. *Neurological Disorders and Pregnancy*. London: Elsevier; 2011:135-158. doi:10.1016/b978-0-12-384911-3.00008-7
 15. Alipannahpour S, Zarshenas M, Ghodrati F, Akbarzadeh M. The severity of post-abortion stress in spontaneous, induced and forensic medical center permitted abortion in Shiraz, Iran, in 2018. *Iran J Nurs Midwifery Res*. 2020;25(1):84-90. doi:10.4103/ijnmr.IJNMR_36_19
 16. Kashfi SM, Yazdankhah M, Babaei Heydarabadi A, Khani Jaihooni A, Tabrizi R. The relationship between religious attitude and mental health in students of Shiraz University of Medical Sciences. *Journal of Research on Religion & Health*. 2016;1(3):34-40. [Persian].
 17. Behere PB, Das A, Yadav R, Behere AP. Religion and mental health. *Indian J Psychiatry*. 2013;55(Suppl 2):S187-194. doi:10.4103/0019-5545.105526
 18. Hamama-Raz Y, Hemmendinger S, Buchbinder E. The unifying difference: dyadic coping with spontaneous abortion among religious Jewish couples. *Qual Health Res*. 2010;20(2):251-261. doi:10.1177/1049732309357054
 19. Akbarzadeh M, Mokhtaryan T, Amooee S, Moshfeghy Z, Zare N. Investigation of the effect of religious doctrines on religious knowledge and attitude and postpartum blues in primiparous women. *Iran J Nurs Midwifery Res*. 2015;20(5):570-576. doi:10.4103/1735-9066.164586
 20. Gholamzadeh S, Godrati F, Saadatmand N, Akbarzadeh M. The obstetrics and gynecology and genetic counseling of mother with legal abortion had been referred to in Fars province center since 2007-2013. *Shiraz E-Med J*. 2016;17(2):e35271. doi:10.17795/semj35271
 21. Godrati F, Saadatmand N, Dinpazhoh M, Akbarzadeh M. Epidemiological study of legal abortion due to fetal defects in the files referred to Fars province forensic medicine centers from 2007 to 2013. *Shiraz E-Med J*. 2016;17(11):e40023. doi:10.17795/semj40023
 22. Ghodrati F, Saadatmand N, Gholamzadeh S, Akbarzadeh M. The seven-year epidemiological study of legal abortion caused by heart disease, blood disorders, diabetes and hypertension as referred to forensic medicine centers in Fars province. *Fam Med Prim Care Rev*. 2019;21(1):23-29. doi:10.5114/fmpcr.2019.82975
 23. Ghodrati F, Saadatmand N, Gholamzadeh S, Akbarzadeh M. Investigation of the prevalence and causes and of legal abortion of teenage married mothers in Iran. *Int J Adolesc Med Health*. 2018;32(1). doi:10.1515/ijamh-2017-0091
 24. Cowchock FS, Lasker JN, Toedter LJ, Skumanich SA, Koenig HG. Religious beliefs affect grieving after pregnancy loss. *J Relig Health*. 2010;49(4):485-497. doi:10.1007/s10943-009-9277-3
 25. Raghobi M, Shirabadi AA, Moallemi S, Narimani M. Demographic characteristics and post-traumatic stress disorder in prison inmates of Zahedan, Iran. *Mil Caring Sci*. 2016;3(1):10-17. doi:10.18869/acadpub.mcs.3.1.10
 26. Bashpour S. The effectiveness of cognitive processing therapy on the improvement of posttraumatic symptoms qol, self esteem and marital satisfaction in the women exposed to infidelity. *Faslnameh Moshavereh va Ravan Darmani Kanevadeh*. 2012;2(2):193-208. [Persian].
 27. Keane TM, Caddell JM, Taylor KL. Mississippi Scale for Combat-Related Posttraumatic Stress Disorder: three studies in reliability and validity. *J Consult Clin Psychol*. 1988;56(1):85-90. doi:10.1037//0022-006x.56.1.85
 28. Goodarzi MA. Evaluating validity and reliability of Mississippi Post Traumatic stress disorder scale. *J Psychol*. 2003;7:153-178.
 29. Alden LE, Regambal MJ, Laposa JM. The effects of direct versus witnessed threat on emergency department healthcare workers: implications for PTSD criterion A. *J Anxiety Disord*. 2008;22(8):1337-1346. doi:10.1016/j.janxdis.2008.01.013
 30. Ebrahimi A, Neshat Doost H, Kalantari M, Molavi H, Asadollahi G. Factor structure, reliability and validity of Religious Attitude Scale. *Journal of Fundamentals of Mental Health*. 2008;10(38):107-116. doi:10.22038/jfmh.2008.1764
 31. Mokhtaryan T, Yazdanpanahi Z, Akbarzadeh M, Amooee S, Zare N. The impact of Islamic religious education on anxiety level in primipara mothers. *J Family Med Prim Care*. 2016;5(2):331-337. doi:10.4103/2249-4863.192314
 32. Allahdadian M, Irajpour A. The role of religious beliefs in pregnancy loss. *J Educ Health Promot*. 2015;4:99. doi:10.4103/2277-9531.171813
 33. Layer SD, Roberts C, Wild K, Walters J. Postabortion grief: evaluating the possible efficacy of a spiritual group intervention. *Res Soc Work Pract*. 2004;14(5):344-350. doi:10.1177/1049731504265829

34. Khazaei H, Rezaei M, Ghadami MR, Tahmasian M, Ghasemi Mobarra A, Shiri E. Relationship between religious values and anxiety among Kermanshah University of Medical Sciences students. *J Kermanshah Univ Med Sci.* 2010;14(1):66-72. [Persian].
35. Bastani F, Ghasemi E, Ramezanzadeh Tabriz E, Janani L, Rahmatnejad L. The investigation of perceived stress and religious coping among female caregivers of the elderly with dementia. *J Rafsanjan Univ Med Sci.* 2015;13(10):925-936. [Persian].
36. Hossieni Garavandi A, Yazdi Ravandi S, Haj Sadeghi Z. Role of androgenic trends, parenting practices, and negative attributional style on spiritual coping among students of Islamic Azad University of Ahvaz, Iran, 2016. *Community Health J.* 2018;11(1):30-39. [Persian].
37. Groleau D, Whitley R, Lespérance F, Kirmayer LJ. Spiritual reconfigurations of self after a myocardial infarction: influence of culture and place. *Health Place.* 2010;16(5):853-860. doi:10.1016/j.healthplace.2010.04.010
38. Nooney JG. Religion, stress, and mental health in adolescence: findings from add health. *Rev Relig Res.* 2005;46(4):341-354. doi:10.2307/3512165
39. Vaillant G, Templeton J, Ardel M, Meyer SE. The natural history of male mental health: health and religious involvement. *Soc Sci Med.* 2008;66(2):221-231. doi:10.1016/j.socscimed.2007.09.011
40. Ghodrati F, Mokhtariyan T, Akbarzadeh M. Islamic-based preventive strategies for postpartum blues. *Sch J Appl Med Sci.* 2016;4(3E):954-958.
41. Tayebi N, Khooshab E, Ghodrati F, Akbarzadeh M. Investigation of correlation between religious attitude and mother-adolescent girls conflict. *J Family Med Prim Care.* 2019 Sep; 8(9): 2893-2897. doi: 10.4103/jfmpc.jfmpc_387_19. PMID: PMC6820374
42. Ghodrati F, Mokhtaryan T, Akbarzadeh M. The effect of pregnancy-related religious training on religious attitudes among pregnant women. *J Midwifery Reprod Health.* 2018;6(3):1305-1313. doi:10.22038/jmrh.2018.23265.1248
43. Geerinck-Vercammen CR, Kanhai HH. Coping with termination of pregnancy for fetal abnormality in a supportive environment. *Obstet Gynecol Surv.* 2004;59(1):16-17. doi:10.1097/01.ogx.0000102784.16345.c4
44. Abboud L, Liamputtong P. When pregnancy fails: coping strategies, support networks and experiences with health care of ethnic women and their partners. *J Reprod Infant Psychol.* 2005;23(1):3-18. doi:10.1080/02646830512331330974
45. Maker C, Ogden J. The miscarriage experience: more than just a trigger to psychological morbidity? *Psychol Health.* 2003;18(3):403-415. doi:10.1080/0887044031000069343
46. Major B, Cozzarelli C, Cooper ML, et al. Psychological responses of women after first-trimester abortion. *Arch Gen Psychiatry.* 2000;57(8):777-784. doi:10.1001/archpsyc.57.8.777
47. Steinberg JR, Tschann JM, Furgerson D, Harper CC. Psychosocial factors and pre-abortion psychological health: the significance of stigma. *Soc Sci Med.* 2016;150:67-75. doi:10.1016/j.socscimed.2015.12.007
48. Biggs MA, Gould H, Foster DG. Understanding why women seek abortions in the US. *BMC Womens Health.* 2013;13:29. doi:10.1186/1472-6874-13-29
49. Foster DG, Steinberg JR, Roberts SC, Neuhaus J, Biggs MA. A comparison of depression and anxiety symptom trajectories between women who had an abortion and women denied one. *Psychol Med.* 2015;45(10):2073-2082. doi:10.1017/s0033291714003213
50. Steinberg JR, Finer LB. Examining the association of abortion history and current mental health: a reanalysis of the National Comorbidity Survey using a common-risk-factors model. *Soc Sci Med.* 2011;72(1):72-82. doi:10.1016/j.socscimed.2010.10.006
51. Steinberg JR, Tschann JM, Henderson JT, Drey EA, Steinauer JE, Harper CC. Psychological distress and post-abortion contraceptive method effectiveness level chosen at an urban clinic. *Contraception.* 2013;88(6):717-724. doi:10.1016/j.contraception.2013.08.009
52. Krok D. Religiousness, spirituality, and coping with stress among late adolescents: a meaning-making perspective. *J Adolesc.* 2015;45:196-203. doi:10.1016/j.adolescence.2015.10.004
53. Becker G, Xander CJ, Blum HE, et al. Do religious or spiritual beliefs influence bereavement? a systematic review. *Palliat Med.* 2007;21(3):207-217. doi:10.1177/0269216307077327
54. García FE, Páez D, Reyes-Reyes A, Álvarez R. Religious coping as moderator of psychological responses to stressful events: a longitudinal study. *Religions.* 2017;8(4):62. doi:10.3390/rel8040062
55. Pereira J, Pires R, Canavarró MC. Psychosocial adjustment after induced abortion and its explanatory factors among adolescent and adult women. *J Reprod Infant Psychol.* 2017;35(2):119-136. doi:10.1080/02646838.2016.1276281
56. Canário C, Figueiredo B, Ricou M. Women and men's psychological adjustment after abortion: a six months prospective pilot study. *J Reprod Infant Psychol.* 2011;29(3):262-275. doi:10.1080/02646838.2011.592974
57. Pope LM, Adler NE, Tschann JM. Postabortion psychological adjustment: are minors at increased risk? *J Adolesc Health.* 2001;29(1):2-11. doi:10.1016/s1054-139x(01)00212-9
58. Frey C. Posttraumatic stress disorder and culture. In: Yilmaz AT, Weiss MG, Riecher-Rössler A, eds. *Cultural Psychiatry: Euro-International Perspectives.* Basel: Karger Publishers; 2001:103-116.
59. Shalev AY, Tuval-Mashiach R, Hadar H. Posttraumatic stress disorder as a result of mass trauma. *J Clin Psychiatry.* 2004;65 Suppl 1:4-10.